

Cushing's Syndrome.

From the dog they thought they'd lost.
A new vitality found.

The timeline of improvement




Efficacy of Vetoryl treatment

According to vets who have treated a number of cases of Cushing's syndrome, owners soon notice the difference and comment on the changes in their pet.

Improvement of clinical signs such as polydipsia, polyuria, polyphagia, panting and lethargy occurs shortly after the start of treatment.

Skin, coat and muscle changes take longer to reverse (usually 3-6 months).

The timeline of improvement



Is your patient responding as expected?


YES

Continue to monitor your patient at regular intervals and encourage its owner to maintain good home records. This will ensure the clinical improvement continues.

NO

If still showing any signs consistent with Cushing's syndrome consider whether a dose increase or an increase in dose frequency would be of benefit to the dog. For further information refer to the treatment & monitoring flowchart.

If your patient is unwell at any time, Vetoryl should be stopped and further investigations undertaken.



Find out more: www.dechra.co.uk/cushings-connect

References

i Internal report VET60

VETORYL: Vetoryl contains trilostane UK: POM-V IE: POM

For further information contact: Dechra Veterinary Products Limited, Sansaw Business Park, Hadnall, Shrewsbury, Shropshire, SY4 4AS
T +44 (0)1939 211200 F +44 (0)1939 211201 www.dechra.co.uk www.dechra.ie
Registered Office: 24 Cheshire Avenue, Cheshire Business Park, Lostock Gralam, Northwich CW9 7UA. Registered in England and Wales, Company Registration No.5385888. Dechra Veterinary Products Limited is a trading business of Dechra Pharmaceuticals PLC